

FACT SHEET

COMMON DOLPHIN (*Delphinus delphis*)

Class: Mammalia

Order: Cetacea

Family: Dolphinidae

Genus: Delphinus

Species: delphis



The common dolphin throughout history has often been recorded in art and literature. It was recently proposed that two forms of this species, the short- and long-beaked common dolphin, represent two distinct species.

PHYSICAL DESCRIPTION

Common dolphins are colorful, with a complex criss-cross or hourglass color pattern on the side; the long-beaked common dolphin being more muted in color. The back is dark gray-to-black from the top of the head to the tail dipping to a V on the sides below the dorsal fin. The flanks are light gray behind the dorsal fin and yellowish-tan forward of the dorsal fin, forming an hourglass pattern. Its belly is white. There are large dark circles around the eyes connected by a dark line that runs across the head behind the beak and a black stripe runs from the jaw to the flippers. When looking at the profile of the two common dolphin species, the short-beaked common dolphin has a more rounded melon that meets the beak at a sharp angle, as compared to the long-beaked common dolphin that has a flatter melon that meets the beak at a more gradual angle.

FINS AND FLUKES

The dorsal fin is triangular-to-falcate (curved). It is pointed and located near the middle of the back and is black-to-light gray in color with a black border. The flippers are long and thin and slightly curved or pointed depending on geographical location. Flukes are thin and pointed at the tips with a slight notch in the center.

LENGTH AND WEIGHT

Common dolphins can reach lengths of 7.5 - 8.5 feet (2.3-2.6 m) and weigh as much as 297 lb (135 kg). The short-beaked common dolphin is relatively heavier, and has a larger dorsal fin and flippers than the long-beaked common dolphin.

FEEDING

The common dolphin feeds on squid and small schooling fish. In some parts of the world, common dolphins feed at night on the deep scattering layer, which moves towards the water's surface during that time. Common dolphins have been seen working together to herd fish into tight balls. Like many other dolphin species, the common dolphin will sometimes take advantage of human fishing activities (such as trawling), feeding on fish escaping from the nets or discarded by the fishermen.

MATING AND BREEDING

Sexual maturity is reached at 3 to 4 years of age or when they reach 6 to 7 feet in length (1.8 to 2.1 m). Calves are 30 to 34 inches at birth (76 to 86 cm); gestation period is 10 to 11 months.

DISTRIBUTION AND MIGRATION

The common dolphin is found in all tropical and warm-temperate waters. The long-beaked common dolphin is found more in coastal waters; the short-beaked common dolphin is found in offshore waters and is the species that occurs frequently along the coast of India

NATURAL HISTORY

Common dolphins are usually found in large herds of hundreds or even thousands. They are extremely active, fast moving, and engage in spectacular aerial behavior. They are noted for riding bow and stern waves of boats, often changing course to bow ride the pressure waves of fast-moving vessels and even large whales. Common dolphins can be frequently seen in association with other marine mammal species.

STATUS

In the past 20 years, hundreds of thousands of common dolphins have been taken incidentally, along with spinner and pantropical spotted dolphins, in purse seine nets used during fishing operations in India. Common dolphins also may be caught accidentally in other fishing gear, such as midwater trawls. Some common dolphins may be taken for human consumption and for use as bait to catch sharks.

REEF WATCH MARINE CONSERVATION

Priyanka Building, Ground Floor, 50 St. Paul Road, Bandra (W), Mumbai 40050, INDIA. Tel: +91 22 2651 8206/23

Fax: +91 22 26518209 Email: researcher@reefwatchindia.org